

to appear in The Astrophysical Journal (July 1, 2000)

One-Line Redshifts and Searches for High-Redshift Ly α Emission¹

Daniel Stern², Andrew Bunker³, Hyron Spinrad
Department of Astronomy, University of California at Berkeley
Berkeley, CA 94720

and

Arjun Dey⁴
KPNO/NOAO
950 N. Cherry Avenue, P.O. Box 26732
Tucson, AZ 85726

ABSTRACT

We report the serendipitous discovery of two objects close in projection with fairly strong emission lines at long wavelength ($\lambda \sim 9190 \text{ \AA}$). One (A) seems not to be hosted by any galaxy brighter than $V_{555} = 27.5$, or $I_{814} = 26.7$ (Vega-based 3σ limits in $1''.0$ diameter apertures), while the other line is associated with a faint ($I_{814} \simeq 24.4$) red galaxy (B) offset by $2''.7$ and $7 \text{ \AA}$ spectrally. Both lines are broad ($\text{FWHM} \approx 700 \text{ km s}^{-1}$), extended spatially, and have high equivalent widths ($W_{\lambda}^{\text{obs}}(\text{A}) > 1225 \text{ \AA}$, 95% confidence limit; $W_{\lambda}^{\text{obs}}(\text{B}) \approx 150 \text{ \AA}$). No secondary spectral features are detected for galaxy A. Blue continuum and the marginal detection of a second weak line in the spectrum of galaxy B is consistent with [O II] (the strong line) and Mg II (the weak line) at $z = 1.466$. By association, galaxy A is likely at $z = 1.464$, implying a rest-frame equivalent width of the [O II] emission line in excess of $600 \text{ \AA}$ and a projected separation of $30 h_{50}^{-1} \text{ kpc}$ for the galaxy pair. Conventional wisdom states that isolated emission lines with rest-frame equivalent widths larger than $\sim 200 \text{ \AA}$ are almost exclusively Ly α . This moderate-redshift discovery therefore compromises recent claims of high-redshift Ly α emitters

¹Based on observations at the W.M. Keck Observatory, which is operated as a scientific partnership among the University of California, the California Institute of Technology, and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W.M. Keck Foundation.

²Current address: Jet Propulsion Laboratory, California Institute of Technology, Mail Stop 169-327, Pasadena, CA 91109; stern@zwoelfkinder.jpl.nasa.gov

³Current address: Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, England

⁴Hubble Fellow

for which other criteria (i.e., line profile, associated continuum decrements) are not reported. We discuss observational tests to distinguish Ly α emitters at high redshift from foreground systems.

Subject headings: galaxies: distances and redshifts – galaxies: evolution – galaxies : formation – early universe

1. Introduction

Serendipitous detections of emission line galaxies are common on low-dispersion spectrograms taken with large ground-based telescopes. Indeed, finding distant galaxies through blank sky slit spectroscopy is fully complementary to narrow-band imaging searches for distant line-emitting galaxies: rather than probing a large area of sky for objects over a limited range of redshift, deep slit spectroscopy surveys a smaller area of sky for objects at a larger range in redshift (e.g., Pritchet 1994; Thompson & Djorgovski 1995). Also, since the resolution of optical spectra is better matched to narrow line emission than filters with widths of $\approx 3000 \text{ km s}^{-1}$, deep slit spectroscopy is vastly more sensitive than narrow-band imaging. In a 1.5 hour spectrum with the Keck telescope at moderate dispersion ($\lambda/\Delta\lambda \simeq 1000$), the limiting flux density probed for spectrally unresolved line emission in a 1 arcsec^2 aperture is $S_{\text{lim}}(3\sigma) \approx 6 \times 10^{-18} \text{ ergs cm}^{-2} \text{ s}^{-1}$ at $\lambda \approx 9300 \text{ \AA}$. The first galaxy confirmed at $z > 5$ was found serendipitously during spectroscopic observations of a galaxy at $z = 4.02$ (Dey et al. 1998). Since then, several other $z > 5$ galaxies have been reported (Hu, Cowie, & McMahon 1998; Weymann et al. 1998; Spinrad et al. 1998; van Breugel et al. 1999; Chen, Lanzetta, & Pascarelle 1999; Hu, McMahon, & Cowie 1999).

In both narrow-band imaging and slit spectroscopy surveys, the equivalent width of detected emission lines is a standard redshift indicator (e.g., Cowie & Hu 1998; Hu et al. 1998). Ly α at high redshift can have a very large rest-frame equivalent width (up to $W_{\lambda}^{\text{rest}} \approx 200 \text{ \AA}$ if driven by star formation; Charlot & Fall 1993), while [O II] $\lambda 3727 \text{ \AA}$ at low to moderate redshift, the other primary strong, solitary emission feature in UV/optical galaxy spectra, rarely has a rest-frame equivalent width exceeding $100 \text{ \AA}$ in magnitude-limited surveys (e.g., Songaila et al. 1994; Guzmàn et al. 1997; Hammer et al. 1997; Hogg et al. 1998). Equivalent width selection is also helped by $W_{\lambda}^{\text{obs}} = W_{\lambda}^{\text{rest}}(1 + z)$, which boosts Ly α more than [O II]. The other strong UV/optical emission features in galaxies, e.g., H β , [O III] $\lambda\lambda 4959, 5007 \text{ \AA}$, H α , are generally easily identified spectroscopically from their wavelength proximity to other emission features, though H α can also have extremely high equivalent widths, up to $3000 \text{ \AA}$ (Leitherer, Carmelle, & Heckman 1995), leading to some confusion between low-redshift, young, dwarf starbursts and high-redshift Ly α -emitters (e.g., Stockton & Ridgway 1998).

Here we report the discovery of two emission lines in the approximate direction of the Abell 2390 cluster. The equivalent widths of these lines are rather large, with $W_{\lambda}^{\text{obs}}(\text{A}) > 1225 \text{ \AA}$ (95%




Fig. 1.— *HST* *I*-band (F814W; 10500 s) image of the serendipitously discovered galaxies in the field of Abell 2390. Galaxy A, marked by dotted circle, is undetected ($I_{814} > 26.7, 3\sigma$). Galaxy B has $I_{814} = 24.4$ ($1''$ diameter aperture). Slit positions for our three observations are indicated. The images shown are $30''$ on a side, oriented with north upwards and east to the left. Galaxy B is located at $\alpha = 21^h53^m35^s.16$, $\delta = +17^\circ42'56''.66$ (J2000) and is $2''.7$ SE of galaxy A. The bright elliptical galaxy south of galaxy B is a member of the Abell 2390 galaxy cluster ($z = 0.241$).

confidence limit) for the first object and $W_\lambda^{\text{obs}}(\text{B}) \approx 150 \text{ \AA}$ for the second object. However, as we argue below, [O II] at $z = 1.46$ is the most likely identification for these features, indicating an atypical system.

In §2 – §4 we discuss the observations, redshift determination, and properties of these galaxies. In §5 we consider the implications of this discovery for narrow-band searches for Ly α emission from distant protogalaxies, including a detailed discussion of the observational criteria useful for distinguishing high-redshift Ly α from foreground emission-line galaxies.

Throughout this paper, unless otherwise indicated, we adopt $H_0 = 50 h_{50} \text{ km s}^{-1} \text{ Mpc}^{-1}$, $q_0 = 0.1$, and $\Lambda = 0$. For these parameters, the luminosity distance, d_L , at $z = 1.46$ is $13.88 h_{50}^{-1} \text{ Gpc}$ and $1''$ subtends $11.1 h_{50}^{-1} \text{ kpc}$.

2. Observations

On UT 1997 July 31, an observation of a pair of lensed arcs close to the core of the rich galaxy cluster Abell 2390 (Frye & Broadhurst 1998; Bunker, Moustakas, & Davis 2000) resulted in the serendipitous detection of a strong emission line at $9185 \text{ \AA}$ approximately $70''$ NNE of the arcs along the long slit spectrogram. These observations were made with the Low Resolution Imaging Spectrometer (LRIS; Oke et al. 1995) at the Cassegrain focus of the Keck II Telescope, using the 400 lines mm^{-1} grating ($\lambda_{\text{blaze}} \approx 8500 \text{ \AA}$). The detector is a Tek 2048² CCD with $24\mu\text{m}$ pixels, corresponding to $0''.212 \text{ pix}^{-1}$. The GG495 filter was used to block second order light and the




Fig. 2.— Two-dimensional, flattened, sky-subtracted spectra near the emission lines from our discovery (July; left) and final (September; right) observations. The abscissa indicates wavelength and the ordinate indicates spatial position. The July data do not sample the entire spatial range. Both data sets have been convolved with a Gaussian of width 1 pixel. Galaxy A is detected in both observations and is the upper emission feature in the September data. Galaxy B is only visible in the September data (lower emission feature). Note the lack of continuum associated with galaxy A in both data sets, while galaxy B has faint continuum blue-ward of the emission line, indicating that the emission line is unlikely to be Ly α . The horizontal feature in the July data set is continuum emission from an Abell 2390 cluster member ($z = 0.241$). The negative horizontal feature in the September data is an artifact of our fringe suppression algorithm (see text); it is the negative of a different Abell 2390 cluster member ($z = 0.232$). Faint negatives of galaxies A and B are also evident.

observations sample $\lambda\lambda 6100 - 9900 \text{ \AA}$. The slit width was $1''$, yielding an effective resolution of $\Delta\lambda_{\text{FWHM}} \approx 8 \text{ \AA}$, and the data were binned by a factor of two spatially during the data acquisition. Reductions were done with IRAF and followed standard spectroscopic procedures. Wavelength calibration was verified against telluric emission lines. The night was photometric with a seeing of $\approx 0''.7$ FWHM, and the data were calibrated using observations of BD+174708 (Massey et al. 1988; Massey & Gronwall 1990) procured on the same night. The total integration time was 3600 s with the slit oriented at a position angle of 23° (see Fig. 1). The emission line source (Fig. 2), which we refer to as galaxy A, was spatially extended by $3''$ in this discovery spectrogram and no associated continuum was detected, implying an equivalent width $W_\lambda^{\text{obs}} > 400 \text{ \AA}$ (95% confidence limit). The wavelength of the emission line does not correspond to any prominent lines at the redshift of Abell 2390 ($z = 0.24$). Throughout we have corrected for foreground Galactic extinction using the dust maps of Schlegel, Finkbeiner, & Davis (1998) which have an optical reddening of $E(B - V) = 0.11$ in the direction of Abell 2390, equivalent to extinctions of $A_{555} = 0.33$ and $A_{814} = 0.20$.

A comparison with archival *Hubble Space Telescope* (HST) Wide Field/Planetary Camera 2

(WFPC2; Trauger et al. 1994) images secured by Fort et al. (HST-GO 5352) finds no obvious optical identification for object A. The *HST* imaging was undertaken on UT 1994 December 10 in the F555W (V_{555}) and F814W (I_{814}) filters. The data comprised four orbits in F555W and five in F814W, with each orbit consisting of a single integration of 2100 s. For the combined F555W image (8400 s), the 1σ limiting magnitude reached in 1 square arcsecond is $V_{555} = 28.4$ mag arcsec $^{-2}$, and for the combined F814W image (10500 s), it is $I_{814} = 27.6$ mag arcsec $^{-2}$ (Vega-based magnitudes are adopted throughout). This is consistent with the predicted sensitivity based on the Poissonian counting statistics and the WF readout noise ($5 e^-$). The 3σ limits on the brightness of the host of galaxy A are $V_{555} > 27.5$ and $I_{814} > 26.7$ for a $1''$ diameter aperture. A faint red galaxy, which we refer to as galaxy B, was found nearby ($\approx 2''.7$ to the SE) but *not* coincident with our estimate of the galaxy A position based on our spectroscopic slit observations (see Fig. 1). Galaxy B is faintly detected in F555W ($V_{555} = 27.63 \pm 0.54$) and has $I_{814} = 24.36 \pm 0.05$, where both magnitudes are quoted for $1''$ diameter apertures.

On UT 1997 August 24 & 26 we observed galaxy B using LRIS in slitmask mode at a position angle of -10.4° (Fig. 1). The $1.4''$ wide slitlet was $42''$ long, and observations were made with both the 600 lines mm $^{-1}$ grating (UT 1997 August 24; 3600 s; $\lambda_{\text{blaze}} = 5000\text{\AA}$; $\Delta\lambda_{\text{FWHM}} \approx 7.5\text{\AA}$; $\lambda\lambda 7050 - 9550\text{\AA}$) and the 300 lines mm $^{-1}$ grating (UT 1997 August 26; 4800 s; $\lambda_{\text{blaze}} = 5000\text{\AA}$; $\Delta\lambda_{\text{FWHM}} \approx 14\text{\AA}$; $\lambda\lambda 3800 - 8800\text{\AA}$). Moderate cirrus affected these observations. Our spectrogram of galaxy B shows a *weak* emission line at $\lambda \simeq 9196\text{\AA}$, nearly the same wavelength as for galaxy A. However, galaxy B also shows weak continuum blue-ward of the emission line extending down to wavelengths of $\sim 4000\text{\AA}$.

Finally, on UT 1997 September 11, we attempted to put the LRIS $1''.0$ wide slit across both galaxies, as well as could be determined from their spectroscopic positions (Fig. 1). These observations were obtained with the 400 lines mm $^{-1}$ grating ($\lambda_{\text{blaze}} = 8500\text{\AA}$) at a position angle of 80° and sample $\lambda\lambda 5950 - 9730\text{\AA}$. The seeing was $\approx 0''.7$. As the night was not photometric, we applied the UT 1997 July 31 sensitivity function, scaled to maintain the same total flux in the galaxy A emission line. The total integration time was 5400 s. In Fig. 2 we present the July and September two-dimensional, flattened, sky-subtracted spectra near the emission lines. In order to suppress the fringing which adversely affects long wavelength ($\lambda \gtrsim 7200\text{\AA}$) optical spectra, we dithered the telescope between individual exposures and subtract temporally adjacent frames prior to co-adding the data sets. This procedure, which is essential for recovering information on faint objects in the telluric OH bands, should not affect the object spectra, but does leave negative holes in the final two-dimensional spectra (see Fig. 2). Fig. 3 presents the extracted, one-dimensional spectra from the September data and Table 1 summarizes the observed properties of galaxies A and B.

3. Redshift Determination

Understanding objects A and B requires first determining their redshifts. Speculations on one-line redshifts are common among faint galaxy observers. Here we have been spared that fate

Table 1. Observed Properties of Galaxies A and B in the Abell 2390 Field.

Parameter	Galaxy A	Galaxy B	
Line ID	[O II]	Mg II	[O II]
λ_{obs} (Å)	9184.8 ± 0.8	$\simeq 6895$	9191.3 ± 1.4
z	1.464	1.463	1.466
f_{line} (10^{-17} ergs cm $^{-2}$ s $^{-1}$)	8.7 ± 0.8	0.9	7.3 ± 0.9
$L_{[\text{OII}]}$ (h_{50}^{-2} 10^{42} ergs s $^{-1}$)	2.02 ± 0.2		1.70 ± 0.2
f_{cont} (10^{-20} ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$)	-2.2 ± 4.2		51.4 ± 4.5
W_{λ}^{obs} (Å)	> 1225	90	$102 - 189$
FWHM (km s $^{-1}$)	627 ± 62	300	785 ± 94
V_{555} (mag)	> 27.5		27.63 ± 0.54
I_{814} (mag)	> 26.7		24.36 ± 0.05

Note. — Magnitudes are in the Vega system and quoted for 1''0 circular apertures. 3σ limits are presented for the magnitude of galaxy A. Spectroscopic measurements derive from UT 1997 September 11 data and have been corrected for Galactic extinction using an optical reddening of $E(B - V) = 0.11$ (Schlegel et al. 1998). The Mg II line is weakly detected. Parameters for the [O II] lines have been measured using the SPECFIT contributed package within IRAF (Kriss 1994) with a flat (in F_{λ}) continuum and a single Gaussian emission line. Equivalent widths have been calculated with a Monte Carlo analysis using the measured line flux and continuum amplitude with errors subject to the constraint that both must be > 0 . We quote the 95% confidence limit for galaxy A and the 95% confidence interval for galaxy B. Line widths (FWHM) are deconvolved by the instrumental resolution (≈ 8 Å).




Fig. 3.— Spectra of galaxies A and B in the Abell 2390 field from the UT 1997 Sep 11 observations, illustrating the high equivalent width emission lines observed at 9185 Å (galaxy A) and 9191 Å (galaxy B). The total exposure time was 5400 s, and the spectrum was extracted using a $1''.5 \times 1''.5$ aperture.

by the presence of the second *weak* emission line in the spectrum of object B at $\lambda \simeq 6895$ Å. The wavelength ratio with the stronger line at $\lambda = 9191$ Å is 1.333, close to the laboratory measured $[\text{O II}]\lambda 3727\text{Å} / \text{Mg II}\lambda\lambda 2796,2803\text{Å} = 1.332$. This implies a redshift $z = 1.466$ for galaxy B with the stronger line identified, as is usually the case, with $[\text{O II}]\lambda 3727\text{Å}$ (hereinafter $[\text{O II}]$). For non-active galaxies it is slightly unusual to observe $\text{Mg II}\lambda\lambda 2796,2803\text{Å}$ (hereinafter Mg II) in emission. For example, Guzmán et al. (1997) report on a spectroscopic study of 51 compact field galaxies in the flanking fields of the Hubble Deep Field. Of the 9 galaxies at redshifts sufficient for Mg II to be sampled by their observations, only 2 (22%) show Mg II in *emission*. Active galaxies, of course, often show Mg II in emission. The velocity offset between the Mg II and $[\text{O II}]$ lines are cause for slight concern, but similar offsets are common in radio galaxy spectra (e.g., Stern et al. 1999b).

As discussed below, the continuum blueward of the long wavelength, isolated line makes alternate redshift identifications unconvincing.

The question then remains: what is the redshift of object A? Previous experience might suggest that isolated optical emission lines with $W_{\lambda}^{\text{obs}} > \text{several} \times 100 \text{ \AA}$ are exclusively identified with $\text{Ly}\alpha$ at high redshift. For example, 0140+326 RD1 at $z = 5.34$ has $W_{\text{Ly}\alpha}^{\text{obs}} = 600 \pm 100 \text{ \AA}$ (Dey et al. 1998) while HDF 4-473.0 at $z = 5.60$ has $W_{\text{Ly}\alpha}^{\text{obs}} \approx 300 \text{ \AA}$ (Weymann et al. 1998). Magnitude-limited redshift surveys find the W_{λ}^{obs} distribution for [O II], typically the primary doppelgänger for high redshift $\text{Ly}\alpha$, rarely has a rest-frame equivalent width exceeding $100 \text{ \AA}$ (e.g., Songaila et al. 1994; Guzmán et al. 1997; Hammer et al. 1997; Hogg et al. 1998). High values of $W_{[\text{OII}]}$ are occasionally seen in AGN, however. The 3rd Cambridge/Molonglo Radio Catalog (3C/MRC) composite radio galaxy spectrum of McCarthy (1993) has $W_{[\text{OII}]} = 128 \text{ \AA}$ while the lower radio power MIT-Green Bank (MG) composite radio galaxy spectrum of Stern et al. (1999b) has $W_{[\text{OII}]} = 142 \text{ \AA}$. Since the surface density of luminous active galaxies is relatively low, our July data suggested at first that the emission line in object A was enticingly identified with $\text{Ly}\alpha$ at the extremely high redshift of $z = 6.55$. Although this interpretation cannot be *completely* ruled out, the robust identification of the emission line in object B with [O II] at $z = 1.466$ coupled with its projected proximity to object A strongly argues that we are witnessing associated galaxies at moderate redshift. The projected separation at $z = 1.466$ is $30 h_{50}^{-1} \text{ kpc}$. The radial velocity difference between the objects is 220 km s^{-1} .

4. Galaxies A and B as Active Galaxies

Associating the [O II] emission in this system with recent star formation activity is likely inappropriate. The [O II] luminosities imply star formation rates of $\approx 90 h_{50}^{-2} M_{\odot} \text{ yr}^{-1}$ (Kennicutt 1992) which seems improbable given the faintness of the hosts. Using the SPECFIT contributed package within IRAF (Kriss 1994) to fit the emission lines with the [O II] $\lambda\lambda 3726, 3729$ doublet, we derive deconvolved circular velocities $v_c \gtrsim 200/\sin i \text{ km s}^{-1}$ for both galaxies. Applying the high-redshift Tully-Fisher relation (Vogt et al. 1997), these line widths suggest apparent I -band magnitudes $I \lesssim 23.4$, much more luminous than the observed galaxies.

A more natural explanation for the spectral character of this system is to associate the line emission from galaxies A and B with active galactic nuclei (AGN). The line widths are consistent with those seen in radio galaxies and Seyfert galaxies: the deconvolved FWHM of galaxies A and B are ≈ 650 and $\approx 800 \text{ km s}^{-1}$ respectively (fit as a single line), while the composite radio galaxy spectra have $\text{FWHM}_{[\text{OII}]} \gtrsim 1000 \text{ km s}^{-1}$ (McCarthy & Lawrence 1999; Stern et al. 1999b). This interpretation also presents a natural explanation for the high equivalent widths, similar to those seen in other active systems.

Comparison with the FIRST radio catalog (Faint Images of the Radio Sky at Twenty-one cm; Becker, White, & Helfand 1995) reveals no radio source within 1 arcminute of either galaxy to a

limiting flux density of $f_{1.4\text{GHz}} \simeq 1 \text{ mJy}$ (5σ). The traditional demarcation between radio-loud and radio-quiet systems is $\log L_{1.4\text{GHz}}(\text{ergs s}^{-1}\text{Hz}^{-1}) = 32.5$. For an emitted luminosity density $L_\nu \propto \nu^\alpha$, this demarcation corresponds to $S_{1.4\text{GHz}} = 1.36h_{50}^2(1+z)^{1+\alpha} \text{ mJy}$ for $z = 1.46$. The FIRST non-detection therefore does not preclude a weak radio-loud source.

5. Tests for Cosmological One-liners

Several programs are currently underway to search for high-redshift primeval galaxies through deep narrow-band imaging (e.g., see Stern & Spinrad 1999). Table 2 lists several recently discovered high-redshift ($z > 5$) sources and includes lower-redshift, strong line-emitters. The previous generation of narrow-band surveys failed to confirm any field Ly α -emitting protogalaxy candidates (Pritchet 1994; Thompson & Djorgovski 1995). Examples of new programs include the Calar Alto Deep Imaging Survey (CADIS, Thommes et al. 1998; Thommes 1999) which uses a Fabry-Pérot interferometer ($S_{\text{lim}}(5\sigma) \approx 3 \times 10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1}$), the Keck-based narrow-band, interference filter imaging program of Cowie, Hu, and McMahon (Cowie & Hu 1998; Hu, Cowie, & McMahon 1999; $S_{\text{lim}}(5\sigma) \approx 1.5 \times 10^{-17} \text{ ergs cm}^{-2} \text{ s}^{-1}$), and serendipitous searches on deep slit spectra (e.g., this paper; Manning et al. 2000).

Selecting objects on the basis of strong line emission may sample a different galaxy population from the traditional magnitude-limited surveys. In particular, emission-line surveys are much more sensitive to active galaxies and objects undergoing massive bursts of star formation. Determining the redshift and physical origin of line emission is challenging; comparison to field surveys selected on the basis of continuum magnitude is perhaps inappropriate. Samples of line-emitting protogalaxy candidates will shortly become available. We therefore present below a timely and detailed discussion of the observational criteria which can be used to distinguish high-redshift Ly α emission from low-redshift interlopers.

5.1. Equivalent Width

The stellar population synthesis models of Charlot & Fall (1993) predict rest-frame Ly α equivalent widths of $50 - 200 \text{ \AA}$ for dust-free young galaxies. For a constant star formation history, the Ly α luminosity and equivalent width are only somewhat dependent on the star formation rate and are greatest at times less than 10 Myr after the onset of the burst. For comparison, the spectral atlas of nearby galaxies by Kennicutt (1992) shows that the rest-frame equivalent width of the H α + [N II] complex rarely exceeds $200 \text{ \AA}$, that of [O III] $\lambda 5007$ rarely exceeds $100 \text{ \AA}$, H β rarely exceeds $30 \text{ \AA}$, and [O II] rarely exceeds $100 \text{ \AA}$. With the $(1+z)$ amplification of observed equivalent widths, emission lines in the optical with measured equivalent widths larger than $\approx 300 \text{ \AA}$ should be almost exclusively identified with Ly α . See Table 2 for a list of the equivalent widths of several recently reported protogalaxy candidates at $z > 5$.

Table 2. Galaxies Reported/Considered at $z > 5$.

Galaxy	z	W_{λ}^{obs} ($\text{\AA}$)	asymmetric line profile	continuum decrement	Reference
STIS parallel gxyA	6.68	≈ 120	...	x	Chen et al. (1999)
SSA22-HCM1	5.74	350	x	x	Hu et al. (1999)
BR 1202–0725 ser	5.64	> 600	...	x	Hu et al. (1998)
HDF 4-473.0	5.60	≈ 300	x	xx	Weymann et al. (1998)
0140+326 RD1	5.34	600 ± 100	x	x	Dey et al. 1998)
HDF 3-951.0	5.34	N/A	N/A	xx	Spinrad et al. (1998)
TN J0924–2201	5.19	≈ 150	1/2	...	van Breugel et al. (1999)
Abell 2390 serA	1.46	> 1225	no	...	this paper
ERO J164502+4626.4	1.44	115	...	x	Dey et al. (1999)
3C212 B08	0.31	≈ 640	...	no	Stockton & Ridgway (1998)

Note. — Symbols indicate reliability of the criterion considered. Ellipses (...) refer to unreported criteria. ‘x’ refers to a positive result and ‘xx’ refers to an extremely positive result — namely, the two $z > 5$ confirmed galaxies in the HDF whose spectral energy distributions are confirmed flat (in f_{ν}) into the near-infrared. The source HDF 3-951.0 lacks Ly α emission; we therefore list some of its criteria not applicable, ‘N/A’. The ‘1/2’ for the high-redshift radio galaxy TN J0924–2201 refers to the enigmatic result that one observation illustrates asymmetric Ly α emission while a second observation does not.

A number of caveats temper sole reliance on equivalent width arguments to discriminate Ly α from foreground emission. First, high-ionization H II dwarf galaxies can have strong H α emission with very weak continuum. Stockton & Ridgway (1998) report an object (3C212 B08) with a single strong emission line at 8567 Å and $W_{\lambda}^{\text{obs}} \approx 640$ Å. They eventually identify the line as H α at $z = 0.305$ due to a secondary feature at 2% of the strong line whose wavelength matches redshifted He I λ 5876 Å. Starburst models (Leitherer et al. 1995) with continuous star formation and a Salpeter initial mass function over the mass range $0.1 - 100 M_{\odot}$ can have H α equivalent widths as high as 3000 Å up to ages of 3 Myr and can remain above ~ 300 Å up to ages of 100 Myr. Furthermore, extreme H II galaxies with very hot ($> 60,000$ K) stars and low metal abundances can have suppressed low-ionization metallic emission lines such as [N II] and [S II], making H α identification difficult (e.g., Tol 1214–277; Fig. 4o in Terlevich et al. 1991). Such galaxies tend to show He I λ 5876 Å in emission.

AGN offer another potential source of ionizing radiation to stimulate line emission: photoionization by a power law continuum emitted near the central engine combined with shock-excited emission can produce very high equivalent width emission. The composite radio galaxy spectrum in McCarthy (1993) shows several lines with rest-frame equivalent widths in excess of 50 Å. As argued earlier, however, the spectral proximity of several features make the most likely identification of an observed, isolated, high-equivalent width emission feature an ambiguous selection between Ly α and [O II]. Composite radio galaxy spectra have $W_{[\text{O II}]} \approx 135$ Å (McCarthy 1993; Stern et al. 1999b). Occasional sources show extremely high equivalent width [O II]. For example, McCarthy (1991) reports [O II] emission from the radio galaxy B3 0903+428 ($z = 0.907$) with $W_{[\text{O II}]} = 251 \pm 55$ Å (rest-frame) and the northern knot in 3C368 has a rest-frame $W_{[\text{O II}]} = 485$ Å (Dey 1999). Luminous, narrow-lined, radio-quiet AGN, the so-called quasar-II population, are another potential active galaxy source of confusion for one-lined sources, though they remain largely unidentified in observational surveys.

5.2. Asymmetric Line Profile

Star formation at low and high redshift is associated with large-scale outflows. Resonant scattering processes in the outflowing gas will trap Ly α photons short-ward of the systemic velocity of the emission and potentially destroy them through dust absorption. This results in a P-Cygni profile for the Ly α line. At the low signal-to-noise ratio observations typically obtained on objects at $z \gtrsim 5$, this causes an asymmetric Ly α line profile with a broad red wing and a steep cut-off on the blue wing. This asymmetric profile is seen in many of the confirmed $z > 5$ Ly α -emitting sources (e.g., Dey et al. 1998) though similar asymmetries might be mimicked by low signal-to-noise ratio observations of [O II] in the low-density limit ($[\text{O II}]\lambda 3726\text{Å} / [\text{O II}]\lambda 3729\text{Å} = 0.7$).

Is an asymmetric line profile a necessary condition for high-redshift Ly α , or merely a sufficient condition? In addition to the local star-forming galaxies with (1) broad damped Ly α absorption centered at the wavelength corresponding to the redshift of the H II emitting gas and (2) galaxies

with Ly α emission marked by blueshifted absorption features, Kunth et al. (1999) notes a third morphology of Ly α line that is occasionally observed in the local Universe: (3) galaxies showing ‘pure’ Ly α emission, i.e., galaxies which show no Ly α absorption whatsoever. Terlevich et al. (1993) present *IUE* spectra of two examples of ‘pure’ emitters: C0840+1201 and T1247–232, both of which are extremely low-metallicity H II galaxies. Thuan & Izotov (1997) present a high signal-to-noise ratio *HST* spectrum of the latter galaxy, noting that with $Z = Z_{\odot}/23$, it is the lowest metallicity local star-forming galaxy showing Ly α in emission. At high signal-to-noise ratio, the emission line shows multiple superposed narrow absorption features, bringing into question the ‘pure’ designation.

Observations of local star-forming galaxies have two implications for studies of high-redshift Ly α emission. First, they provide a natural explanation for asymmetric profiles which seem to characterize high-redshift Ly α , but also imply that although the asymmetric profile may be a sufficient condition for identification of a strong line with Ly α , it is not a necessary one. Second, if Ly α emission is primarily a function of kinematics and perhaps evolutionary phase of a starburst as suggested by the scenario of Tenorio-Tagle et al. (1999), attempts to derive the comoving star-formation rate at high redshifts from Ly α emission will require substantial and uncertain assumptions regarding the relation of observed Ly α properties to the intrinsic star-formation rate.

5.3. Continuum Decrements

Actively star-forming galaxies should, in the absence of dust absorption, have blue continua, nearly flat in f_{ν} , at rest-frame ultraviolet wavelengths long-ward of Ly α . This radiation derives from the hot, massive, short-lived stars and can therefore be used as indicator of the instantaneous star formation rate of a galaxy modulo uncertainties in dust absorption, age, metallicity, and stellar initial mass function. Short-ward of the Lyman limit (912 Å), the spectral energy distributions should drop steeply. This is due both to photospheric absorption in the UV-emitting stars themselves, as well as photoelectric absorption by neutral hydrogen along the line of sight to the galaxy. Between the Lyman limit and Ly α , photoelectric absorption from neutral hydrogen in the intergalactic medium attenuates the emitted spectrum.

In terms of emission line surveys, this discontinuity serves as a useful foil for identifying observed lines with high-redshift Ly α , provided the data are sufficiently sensitive to detect continuum. Objects which show evidence of a flat spectral energy distribution across an emission line can immediately be ruled out as distant galaxy candidates. The presence of a discontinuity, however, is *not sufficient* for classifying the observed emission line with Ly α , unless the amplitude is extreme (see below). In particular, [O II], the other strong, solitary emission feature in the UV/optical spectra of star-forming galaxies, lies short-ward of the Balmer and 4000 Å breaks. The former is strongest in young systems dominated by an A-star population, while the latter arises from metal-line blanketing (predominantly Fe II) in late-type stars. At high signal-to-noise ratio, the morphology of the break can be used to distinguish the Ly α forest from the Balmer break from the 4000 Å break.

At low signal-to-noise ratio, however, the continuum decrement across [O II] can mimic that across high-redshift Ly α . For example, Dey et al. (1999) report an optical spectrogram of the extremely red object ERO J164502+4626.4 (HR10 in Hu & Ridgway 1994), showing a single, strong emission feature at 9090.6 Å and a drop in the continuum level by a factor of ≈ 3 across the emission line. The optical data alone is suggestive of Ly α emission at $z = 6.48$. However, near-infrared images show the source to be extremely red, with a steeper spectral energy distribution than expected for a high-redshift, star-forming galaxy. Furthermore, a near-infrared spectrum shows a second emission at 1.603 μ m, solidifying the emission line identifications as [O II] and H α at $z = 1.44$. Extremely dusty, moderate-redshift galaxies such as ERO J164502+4626.4 may be misidentified as high-redshift Ly α -emitters by *solely* optical surveys.

The amplitude of a continuum discontinuity may be used as a tool for distinguishing the identification of that discontinuity. In order of decreasing wavelength, discontinuities are commonly observed in UV/optical spectra of galaxies at rest wavelengths of 4000 Å [$D(4000)$], 2900 Å [$B(2900)$], 2640 Å [$B(2640)$], 1216 Å (Ly α), and 912 Å (the Lyman limit). The hydrogen discontinuities derive from associated and foreground absorption and thus have no theoretical maximum. The longer rest-wavelength discontinuities derive from metal absorption in the stars and galaxies and are thus dependent on the age and metallicity of the galaxy (c.f., Fanelli et al. 1992). The largest measured values of $D(4000)$ are ~ 2.6 (Hamilton 1985; Dressler & Gunn 1990), while *IUE* spectra of main-sequence stars exhibit $B(2900) \lesssim 3$ and $B(2640) \lesssim 3$ (Spinrad et al. 1997). The example of ERO J164502+4626.4 illustrates the utility of break amplitudes for redshift identifications. The amplitude of the continuum break in ERO J164502+4626.4 is ≈ 3 across an emission line at 9090.6 Å. Were the emission line Ly α , the implied redshift would be $z = 6.48$. Models and measurements of the strength of the Ly α forest imply decrements $\gtrsim 10$ at $z \gtrsim 6$ (Madau 1995; Stern & Spinrad 1999), incompatible with the observed decrement and arguing for a lower redshift line identification.

6. Conclusions

We report the serendipitous discovery of two faint galaxies with high equivalent width emission lines at long wavelength ($\lambda \approx 9190$ Å). For one source, galaxy B, faint blue continuum and a weak secondary line emission feature implies the source is at $z = 1.466$. The spatial proximity and similar emission wavelength of object A persuasively argues that this is an unusual [O II]-emitter at $z = 1.464$. It had been thought that serendipitous one-lined sources with rest-frame equivalent widths larger than a few $\times 100$ Å are exclusively identified with high-redshift Ly α . Our observations have shown that this is demonstrably not the case. Both sources are unlike local star-forming galaxies, and we suggest that the observations are most consistent with the discovery of a moderate-redshift active system.

Several programs are currently underway to find high-redshift Ly α emitters using a combination of narrow- and broad-band imaging (e.g., Thommes et al. 1998; Hu et al. 1998) and

serendipitous long slit surveys (e.g., Manning et al. 2000). The moderate-redshift system discussed herein has serious implications for those surveys. In particular we expect the line-emission surveys to uncover a population of strong line emitters and we emphasize that comparison of this sample with magnitude-limited surveys could be misleading.

We discuss various criteria which can be used to assess whether a solitary, high-equivalent width emission feature is associated with high-redshift $\text{Ly}\alpha$. In Table 2 we consider how recently reported $z > 5$ spectroscopic candidates fare with respect to these criteria. Some sources are reliably confirmed at $z > 5$, while others are confirmed at low-redshift and yet others remain ambiguous with the current data. We conclude that some criteria are necessary, but not sufficient, to conclude that a source is at $z > 5$, such as a continuum decrement across the emission feature, while other criteria are sufficient, but not necessary, such as an asymmetric line profile. We suggest that multiple criteria are necessary to convincingly demonstrate that a single-lined source is at high-redshift.

We thank Alex Filippenko, Doug Leonard, and Aaron Barth for obtaining the July 1997 observations, and Tom Broadhurst and Brenda Frye for the follow-up observations during August 1997. We are indebted to the expertise of the staff of Keck Observatory for their help in obtaining the data presented herein, and to the efforts of Bev Oke and Judy Cohen in designing, building, and supporting LRIS. The work presented here has been aided by discussions with Mike Liu, Curtis Manning, Gordon Squires, and Chuck Steidel. We are also grateful to Trinh Thuan for sharing the *HST*/GHRS spectrum of T1214–277 and to Carlos De Breuck and Adam Stanford for carefully reading the manuscript. This work has been supported by the following grants: IGPP/LLNL 98-AP017 (DS), NICMOS/IDT grant NAG 5-3042 (AJB), NSF grant AST 95-28536 (HS), and NASA HF-01089.01-97A (AD). A preliminary version of this work was presented at the January 1999 AAS meeting in Austin under the title “Cosmological One-Liners: A Cautionary Tale” (Stern et al. 1999a).

REFERENCES

- Becker, R. H., White, R. L., & Helfand, D. J. 1995, *ApJ*, 450, 559
- Bunker, A. J., Moustakas, L. A., & Davis, M. 2000, *ApJ*, in press (March)
- Charlot, S. & Fall, S. M. 1993, *ApJ*, 378, 471
- Chen, H.-W., Lanzetta, K. M., & Pascarelle, S. 1999, *Nature*, 398, 586
- Cowie, L. & Hu, E. M. 1998, *AJ*, 115, 1319
- Dey, A. 1999, in *The Most Disant Radio Galaxies*, ed. H. Röttgering, P. N. Best, & M. D. Lehnert (Dordrecht: Kluwer), 19

- Dey, A., Graham, J. R., Ivison, R. J., Smail, I., Wright, G. S., & Liu, M. C. 1999, *ApJ*, 519, 610
- Dey, A., Spinrad, H., Stern, D., Graham, J. R., & Chaffee, F. 1998, *ApJ*, 498, L93
- Dressler, A. & Gunn, J. E. 1990, in *Evolution of the Universe of Galaxies: the Edwin Hubble Centennial Symposium*, ed. R. G. Kron, Vol. 10 (San Francisco: ASP Conference Series), 200
- Fanelli, M. N., O’Connell, R. W., Burstein, D., & Wu, C. C. 1992, *ApJS*, 82, 197
- Frye, B. & Broadhurst, T. J. 1998, *ApJ*, 499, 115
- Guzmán, R. et al. 1997, *ApJ*, 489, 559
- Hamilton, D. 1985, *ApJ*, 297, 371
- Hammer, F. et al. 1997, *ApJ*, 481, 49
- Hogg, D. W. et al. 1998, *ApJ*, 504, 622
- Hu, E. M., Cowie, L. L., & McMahon, R. G. 1998, *ApJ*, 502, 99
- Hu, E. M., McMahon, R. G., & Cowie, L. L. 1999, *ApJ*, 522, 9
- Hu, E. M. & Ridgway, S. E. 1994, *AJ*, 107, 1303
- Kennicutt, R. 1992, *ApJ*, 388, 310
- Kriss, G. 1994, in *Astronomical Data Analysis Software and Systems III*, Vol. 61 (San Francisco: ASP Conference Series), 437
- Kunth, D., Terlevich, E., Terlevich, R., & Tenorio-Tagle, G. 1999, in *Dwarf Galaxies and Cosmology*, ed. T. X. Thuan (Gif-sur-Yvette: Editions Frontieres), in press
- Leitherer, C., Carmelle, R., & Heckman, T. M. 1995, *ApJS*, 99, 173
- Madau, P. 1995, *ApJ*, 441, 18
- Manning, C., Stern, D., Spinrad, H., & Bunker, A. J. 2000, *ApJ*, submitted
- Massey, P. & Gronwall, C. 1990, *ApJ*, 358, 344
- Massey, P., Strobel, K., Barnes, J. V., & Anderson, E. 1988, *ApJ*, 328, 315
- McCarthy, P. J. 1991, *AJ*, 102, 518
- . 1993, *ARA&A*, 31, 639
- McCarthy, P. J. & Lawrence, C. R. 1999, *ApJ*, in preparation

- Oke, J. B. et al. 1995, PASP, 107, 375
- Pritchett, C. J. 1994, PASP, 106, 1052
- Schlegel, D., Finkbeiner, D., & Davis, M. 1998, ApJ, 500, 525
- Songaila, A., Cowie, L. L., Hu, E. M., & Gardner, J. P. 1994, ApJS, 94, 461
- Spinrad, H., Dey, A., Stern, D., Peacock, J. A., Dunlop, J., Jimenez, R., & Windhorst, R. A. 1997, ApJ, 484, 581
- Spinrad, H., Stern, D., Bunker, A. J., Dey, A., Lanzetta, K., Yahil, A., Pascarelle, S., & Fernández-Soto, A. 1998, AJ, 116, 2617
- Stern, D., Bunker, A. J., Spinrad, H., & Dey, A. 1999a, BAAS, 193, #122.09
- Stern, D., Dey, A., Spinrad, H., Maxfield, L. M., Dickinson, M. E., Schlegel, D., & González, R. A. 1999b, AJ, 117, 1122
- Stern, D. & Spinrad, H. 1999, PASP, 111, 1475
- Stockton, A. & Ridgway, S. E. 1998, AJ, 115, 1340
- Tenorio-Tagle, G., Kunth, D., Terlevich, E., Terlevich, R., & Silich, S. A. 1999, MNRAS, in press, astro-ph/9905324
- Terlevich, E., Díaz, A. I., Terlevich, R., & García, M. L. 1993, MNRAS, 260, 3
- Terlevich, R., Melnick, J., Masegosa, J., Moles, M., & Copetti, M. V. F. 1991, A&AS, 91, 285
- Thommes, E. 1999, in *From Stars to Galaxies to the Universe*, in press, astro-ph/9812223
- Thommes, E., Meisenheimer, K., Fockenbrock, R., Hippelein, H., Röser, H. J., & Beckwith, S. 1998, MNRAS, 293, L6
- Thompson, D. & Djorgovski, S. G. 1995, AJ, 110, 982
- Thuan, T. X. & Izotov, Y. I. 1997, ApJ, 489, 623
- Trauger, S. et al. 1994, ApJ, 435, 3
- van Breugel, W., De Breuck, C., Stanford, S. A., Stern, D., Röttgering, H., & Miley, G. 1999, ApJ, 518, L61
- Vogt, N. et al. 1997, ApJ, 435, 3
- Weymann, R., Stern, D., Bunker, A. J., Spinrad, H., Chaffee, F., Thompson, R., & Storrie-Lombardi, L. 1998, ApJ, 505, L95